

KINGDOM OF CAMBODIA
National Religion King



3

Ministry of Agriculture Forestry and Fisheries
Fisheries Administration

3

**Cambodia Programme for Sustainable and Inclusive Growth
in the Fisheries Sector: Capture Component**

Monthly Statistical Report
Scientific Catch Assessment of Inland Fisheries in Cambodia
July 2022

By Inland Fisheries Research and Development Institute

Funded by European Union
ACA/2018/041-466 and ACA/2019/041-594

1. Introduction

IFReDI, with technical assistance from FAO CAPFISH project under EU budget support, is implementing a scientific catch assessment survey, using a monthly household recall survey for inland fisheries in Cambodia. The aim is to obtain better information on catch and effort by small-scale household fisheries in Cambodia, and to develop a sustainable catch monitoring methodology for implementation by provincial fisheries administrations, supported by IFReDI. The current statistical report provides preliminary analysis based on the available data and focuses on the main indicators that are covered by the catch assessment survey. Therefore, the results do not represent final estimates and may be changed in future updates.

Methodology of data collection and analysis

A description of the methodology can be found in: Fisheries Administration (FiA). 2021. Manual for Scientific Catch Assessment by Recall survey of Inland Fisheries in Cambodia. Inland Fisheries Research and Development Institute of the Fisheries Administration, Phnom Penh, Cambodia. 47 pages.



The total estimated catch in this report is calculated using the proportion of fishing households found by the random household selection under the Household Selection Interview survey. This is taken to be representative for the proportion of fishing households for each fishing area and this is combined with the total number of rural households by fishing area from the NIS 2019 population census to estimate the total number of fishing households. The Fishing Activity Coefficient is estimated from proportion of households reporting fishing activities in the Household Catch Interview.

Estimates for CPUE, the average (mean) daily household catch and the mean monthly household catch used for extrapolating the total catch, come with a value for the relative standard error ($\epsilon\%$). This is used to indicate the accuracy of the estimate for the mean catch. If the $\epsilon\%$ is higher than 30% this indicates a high inaccuracy¹, due to high variation or low sample size and the resulting total estimate should not be used.

In tables with the proportion of reported catch obtained by habitat and fishing gear, the average daily catch by habitat or gear (CPUE) isn't included. The available data displays too much variation and the accuracy is too low for it to be statistically accurate and representative for the real CPUE. This will be included, where relevant in the annual report.

¹ For national statistical reports the rule of thumb states that if the *relative* standard error is higher than 30%, the average should not be reported. The current report includes all estimated values to indicate that an estimate is available, with the $\epsilon\%$ indicating the statistical accuracy.

3. Statistical tables and results

Table 1. Number of random selected households covered by the survey and proportion of target household by fishing area for July 2022.

Fishing Area	Villages	Households		
		Count	Target	Proportion
Floodplain	8	120	120	100.0%
Plateau	8	102	120	85.0%
Tonle Sap	10	142	150	94.7%
Total	26	364	390	93.3%

Table 2. Mean **daily** household catch (CPUE), with number of active fishing households, standard deviation and relative standard error, by fishing area.

Fishing Area	Active HH	Daily HH catch (Kg)	SD	ε%
Floodplain	46	2.36	4.23	26.4%
Plateau	63	2.55	3.46	17.1%
Tonle Sap	92	2.63	2.73	10.9%
Grand Total	201	2.54	3.34	9.3%

Mean daily catch calculated based on the reported 5-day catch and fishing days, with SD is Standard Deviation; ε% is relative Standard Error

Table 3. Mean **monthly** household catch, with proportion of active fishing households, standard deviation, relative standard error and total estimated catch by fishing area.

Fishing Area	% Active HH	Monthly HH catch (Kg)	SD	ε%	Total (MT)
Floodplain	38%	30.37	6.28	18.3%	7,453
Plateau	62%	57.01	12.25	16.2%	3,434
Tonle Sap	65%	33.66	8.13	15.1%	8,302
Grand total	55%	40.22	9.44	9.9%	
Total estimated catch (MT)					19,189

SD is Standard Deviation; ε% is relative Standard Error

Table 4. Proportion of fishing days on which male and female adults and children are reporting fishing activities.

Fishing Area	Adult Female	Adult Male	Child Female	Child Male
Floodplain	8.8%	96.5%	4.4%	7.0%
Plateau	5.6%	93.1%	0.0%	7.9%
Tonle Sap	7.8%	98.4%	3.6%	1.6%
Grand Total	7.1%	95.8%	2.3%	5.4%

The maximum involvement of each gender and age group is 100% for each fishing area, if they are fishing on all reported fishing days, the total for each fishing area can be more than 100%.

Table 5. Reported catch (Kg) with proportion caught by main boat type by fishing area.

Fishing Area	Catch (Kg)	No boat	Motorised	Non-motorised
Floodplain	185.6	39.3%	27.3%	33.5%
Plateau	478.5	23.3%	62.3%	14.4%
Tonle Sap	395.8	62.9%	17.5%	19.6%
Grand Total	1,059.9	40.9%	39.4%	19.7%

Overall proportion based on weighted average catch by main boat type and fishing area, not reported total catch²




Figure 1. Overall contribution of the main boat types to total reported catch.

Table 6. Proportion and reported catch by habitat for single habitat catches by fishing area.

Fishing Habitats	Floodplain	Plateau	Tonle Sap	Grand Total
Mekong Mainstream	21.7%	74.3%	0.0%	39.4%
Sub-Stream	9.5%	16.9%	8.4%	12.7%
Floodplain: rice fields (rain)	38.9%	5.8%	9.3%	12.7%
Floodplain: lakes and ponds	9.9%	0.6%	29.5%	12.3%
Tributaries to Tonle Sap	1.1%	0.0%	30.8%	10.9%
Irrigation canals	11.9%	0.0%	8.3%	4.9%
Stream	0.0%	0.0%	11.8%	4.1%
Floodplain: flooded forest	3.1%	2.0%	0.0%	1.5%
Major Tributaries	3.7%	0.0%	0.0%	0.6%
Reservoir	0.0%	0.0%	1.6%	0.6%
Unspecified	0.0%	0.3%	0.3%	0.2%
Seasonal swamps	0.0%	0.1%	0.0%	0.0%
Grand Total (kg)	198.0	556.7	404.1	1,158.8

Only catch for fishing days that report fishing in a single habitat is included

² This is the standard way to calculate, but isn't done for habitat and gear catch, as this is complicated by fishing days where the reported catch is from multiple habitats or caught by multiple gears




Figure 2. Overall contribution of habitats to total reported catch, with proportion of catch for floodplain habitats.

Table 7. Proportion and reported catch by gear for single gear days, by fishing area.

Fishing habitats	Floodplain	Plateau	Tonle Sap	Grand Total
Stationary gillnet	30.4%	46.6%	33.3%	38.9%
Horizontal cylinder trap (small)	14.4%	19.2%	16.0%	17.1%
Cast net	5.9%	3.2%	21.8%	10.3%
Unspecified	1.3%	16.2%	3.6%	9.0%
Drifting gillnet	12.9%	11.0%	0.4%	7.6%
Hook long line	0.0%	2.7%	5.4%	3.1%
Pumping	15.5%	0.0%	0.0%	2.9%
Hand capture	9.4%	0.0%	2.8%	2.7%
Bamboo vertical cylinder trap	1.4%	0.0%	5.9%	2.4%
Horizontal cylinder trap (large)	4.5%	0.0%	4.3%	2.3%
Pole and line	0.0%	0.6%	5.6%	2.2%
Hook and line	3.0%	0.3%	0.0%	0.7%
Spear	1.4%	0.2%	0.0%	0.4%
scoop nets	0.0%	0.0%	1.0%	0.4%
Grand Total (kg)	181.1	444.1	342.31	967.5

Only catch for fishing days that report fishing with a single gear is included, therefore the total is different from reported catch by habitat.




Figure 3. Overall contribution of fishing gears to total reported catch.

Table 8. Reported disposal by fishing area in weight and proportion.

Fishing Area	Sold Kg	% Sold	Consumed Kg	% Consumed	Other use Kg	% Other use
Floodplain	105.3	22.0%	91.6	16.1%	35.9	12.0%
Plateau	159.0	33.2%	225.6	39.6%	214.0	71.4%
Tonle Sap	214.0	44.7%	252.3	44.3%	49.8	16.6%
Grand Total	478.3	35.5%	569.5	42.3%	299.71	22.2%




Figure 4. Catch contribution for fish and other aquatic animals.

Table 9. Top 15 reported species and species groups **by weight** in reported household catch, with reported weight and proportion of catch by individual species and species groups.

No.	Scientific name	Khmer name	Catch (kg)	Catch contribution	
				%Catch	%Cum.
1	<i>Henicorhynchus siamensis</i>	ត្រីរៀលតូប	158.0	11.7%	11.7%
2	<i>Puntioplites proctozysron</i>	ត្រីប្រកែង	113.05	8.4%	20.1%
3	<i>Anabas testudineus</i>	ត្រីក្រញូ	107.4	8.0%	28.1%
4	<i>Henicorhynchus lobatus</i>	ត្រីរៀលអង្កាម	100.3	7.4%	35.5%
5	<i>Channa striata</i>	ត្រីវល្ល/ធូក	84.3	6.3%	41.8%
6	Mixed small or juvenile fish	ត្រីល្អិតចម្រុះ	59.9	4.4%	46.2%
7	<i>Somanniathelphusa sp.</i>	ក្ដាមស្រែ	51.4	3.8%	50.0%
8	<i>Trichopodus trichopterus</i>	ត្រីកំភ្លាញស្រែ	47.4	3.5%	53.6%
9	<i>Cyclocheilichthys enoplos</i>	ត្រីឆ្កោក	37.4	2.8%	56.3%
10	<i>Labiobarbus siamensis</i>	ត្រីអាចម៍កុក	30.5	2.3%	58.6%
11	<i>Hemibagrus wyckioides</i>	ត្រីខ្សា	28.3	2.1%	60.7%
12	<i>Hemibagrus spilopterus</i>	ត្រីឆ្កាំង	26.8	2.0%	62.7%
13	Small mixed shrimps	កំពើសចម្រុះ	26.4	2.0%	64.6%
14	<i>Osteochilus lini</i>	ត្រីក្រស	25.8	1.9%	66.6%
15	<i>Hypsibarbus suvattii</i>	ត្រីឆ្អិនស្ទើង	24.8	1.8%	68.4%
16	Other	ផ្សេងទៀត	425.8	31.6%	100.0%
Total reported catch			1,347.5		

Table 10. Top 15 reported species by value (1000 Riel) in reported catch, with reported value, proportion of value by individual species and species groups.

No.	Scientific name	Khmer name	Value (1000 Riel)	Value Contribution	
				%Value	%Cum.
1	<i>Channa striata</i>	ត្រីវែង/ផ្នក់	1,686	16.8%	16.8%
2	<i>Henicorhynchus siamensis</i>	ត្រីវៀលតុប	1,264	12.6%	29.4%
3	<i>Hemibagrus wyckioides</i>	ត្រីខ្សា	679	6.8%	36.2%
4	<i>Puntioplites proctozysron</i>	ត្រីប្រកែង	622	6.2%	42.4%
5	<i>Anabas testudineus</i>	ត្រីក្រញូ	591	5.9%	48.3%
6	<i>Hemibagrus spilopterus</i>	ត្រីឆ្មាំង	483	4.8%	53.1%
7	<i>Henicorhynchus lobatus</i>	ត្រីវៀលអង្កាម	351	3.5%	56.6%
8	<i>Cyclocheilichthys enoplos</i>	ត្រីឆ្កែក	337	3.4%	60.0%
9	<i>Trichopodus trichopterus</i>	ត្រីកំភ្លាញស្រែ	261	2.6%	62.6%
10	<i>Labiobarbus siamensis</i>	ត្រីអាចម៍កុក	229	2.3%	64.9%
11	Mixed small or juvenile fish	ត្រីល្អិតចម្រុះ	210	2.1%	67.0%
12	<i>Osteochilus lini</i>	ត្រីក្រស	206	2.1%	69.0%
13	<i>Paralaubuca typus</i>	ត្រីស្លឹកឫស្សីធំ	194	1.9%	71.0%
14	<i>Clarias batrachus</i>	ត្រីអណ្តែងរឹង	188	1.9%	72.9%
15	<i>Hypsibarbus suvattii</i>	ត្រីឆ្មិនស្តើង	186	1.9%	74.7%
16	Others		2,534	25.3%	100.0%
Total report value			10,020		

ថ្ងៃ ខែ ឆ្នាំ ព.ស.២៥៦៦
រាជធានីភ្នំពេញ ថ្ងៃទី ខែ ឆ្នាំ ២០២៣
ហត្ថលេខា

បានឃើញ
ប្រតិភូរាជរដ្ឋាភិបាល ទទួលបន្ទុកជា
ប្រធានរដ្ឋបាលជលផល និងជាប្រធានកម្មវិធី **CAPFISH-Capture**
ថ្ងៃ ខែ ឆ្នាំ ព.ស.២៥៦៦
រាជធានីភ្នំពេញ ថ្ងៃទី ខែ ឆ្នាំ ២០២៣

បណ្ឌិត ហេង គង់